arXiv:1209.1945v1 [gr-qc] 10 Sep 2012

Lense-Thirring Precession in Plebański-Demiański spacetimes

Chandrachur Chakraborty*
Saha Institute of Nuclear Physics, Kolkata 700064, India

Partha Pratim Pradhan†
Vivekananda Satabarshiki Mahavidyalaya, Paschim Medinipur, WB 721513, India.

An exact expression of Lense-Thirring precession rate is derived for non-extremal and extremal Plebański-Demiański(PD) spacetimes. This formula is used to find the exact Lense-Thirring precession rate in various axisymmetric spacetimes, like: Kerr[1], Kerr-Newman, Kerr-de Sitter, Kerr-Taub-NUT etc. For the extremal case, we have also computed the general extremality condition for PD spacetimes which could be applied to get the extremal limit in any stationary and axisymmetric spacetimes.

I. INTRODUCTION

General relativity predicts that curvature is produced, not only by the distribution of mass-energy, but also by its motion. This is not unlike electromagnetism, where fields are produced not only by charge distributions but also by currents. Pursuing this analogy, these effects are sometimes called gravitomagnetic. An observer riding with a gyroscope will see its spin precess in any stationary or static spacetime due to the distribution of mass-energy. This effect is called geodetic precession. But, a stationary spacetimes with angular momentum (rotation) are known to exhibit an another effect called Lense-Thirring (LT) precession whereby locally inertial frames are dragged along the rotating spacetime, making any test gyroscope in such spacetimes *precess* with a certain frequency called the LT precession frequency [2]. A gyroscope in such a spacetime undergoes a precession due to this inertial frame-dragging effect. The axisymmetric vacuum solution of the Einstein equation is first derived[3] by R. Kerr. The Kerr solution is then generalized as Kerr-Newman, Kerr-de Sitter, Kerr-Taub-NUT spacetimes etc. The generalization of these axisymmetric vacuum solutions can be expressed as the special cases of Plebański-Demiański (PD) spacetime[4]. In this paper, our aim is to derive the exact precession rate of LT precession in the most general axially symmetric spacetimes, i.e, PD spacetimes. From that general expression of LT precession rate in PD spacetimes, we can easily derive the LT precession rate in Kerr, Kerr-Newman, Kerr-Taub-NUT spacetimes etc. in both for non-extremal and extremal cases. In section II, we shall calculate the LT precession rate for non-extremal axially symmetric spacetimes and in section III, we shall do the same for the extremal spacetimes.

II. NON-EXTREMAL CASE

A. Plebański-Demiański (PD) spacetimes

The PD spacetime is the most general axially symmetric vacuum solution of Einstein equation, at present. The PD spacetimes are characterized by the seven parameters mass (M), Kerr parameter (a, angular momentum per unit mass), acceleration, NUT parameter (n), electric charge (Q_c), magnetic charge (Q_m) and Cosmological constant. The line element of six parameters (as acceleration vanishes) PD spacetimes can be written as (taking, $G = c = 1$),

$$ds^2 = -\frac{\Delta}{p^2}(dt - A d\phi)^2 + \frac{p^2}{\Delta}dr^2 + \frac{p^2}{\Xi}d\theta^2 + \frac{\Xi}{p^2}\sin^2\theta(a dt - B d\phi)^2 \quad (1)$$

where,

$$\begin{aligned} p^2 &= r^2 + (n - a\cos\theta)^2, \\ A &= a\sin^2\theta + 2n\cos\theta, B = r^2 + a^2 + n^2 \\ \Delta &= (r^2 + a^2 - n^2)(1 - \frac{1}{\ell^2}(r^2 + 3n^2)) \\ &\quad - 2Mr + Q_c^2 + Q_m^2 - \frac{4n^2r^2}{\ell^2} \\ \Xi &= 1 + \frac{a^2\cos^2\theta}{\ell^2} - \frac{4an\cos\theta}{\ell^2} \end{aligned} \quad (2)$$

$\frac{1}{\ell^2} = \Lambda$ denotes the Cosmological constant divided by three, represents the PD("de-Sitter") spacetimes and if ℓ^2 is replaced by $-\ell^2$, it represents the PD("AdS") spacetimes. [1] So, our metric $g_{\mu\nu}$ is thus following

---

* chandrachur.chakraborty@saha.ac.in
† pppradhan77@gmail.com

[1] Actually, there are two cases in PD spacetimes, one is for Λ positive which we call as PD("dS") and another is for Λ negative which we can call as PD("AdS").

$$g_{\mu\nu} = \begin{pmatrix} -\frac{1}{p^2}(\Delta - a^2\Xi\sin^2\theta) & 0 & 0 & \frac{1}{p^2}(A\Delta - aB\Xi\sin^2\theta) \\ 0 & \frac{p^2}{\Delta} & 0 & 0 \\ 0 & 0 & \frac{p^2}{\Xi} & 0 \\ \frac{1}{p^2}(A\Delta - aB\Xi\sin^2\theta) & 0 & 0 & \frac{1}{p^2}(-A^2\Delta + B^2\Xi\sin^2\theta) \end{pmatrix} \tag{3}$$

In our previous paper[1], we have already given an exact expression of LT precession rate (eqn.(4) of [1]), which is applicable for any non-accelerating stationary spacetimes. In this paper, we are going to derive the LT precession rate for non-accelerating PD spacetimes and also some others axially symmetric spacetimes like this. The expression for LT precession rate in non-accelerating, stationary and axisymmetric spacetime can be written as,

$$\Omega_{LT} = \frac{1}{2\sqrt{-g}}\left[\left(g_{0\phi,r} - \frac{g_{0\phi}}{g_{00}}g_{00,r}\right)\partial_\theta - \left(g_{0\phi,\theta} - \frac{g_{0\phi}}{g_{00}}g_{00,\theta}\right)\partial_r\right] \tag{4}$$

which is same as eqn.(7) of [1]. The various metric components can be read off from above metric (3). Likewise,

$$\sqrt{-g} = p^2\sin\theta \tag{5}$$

Substituting the metric components into eqn.(4) we can easily get the LT precession rate in PD spacetimes. But, there is a problem in that formulation. So, we should transform the precession frequency formula from the coordinate basis to the orthonormal 'Copernican' basis: first note that

$$\Omega_{LT} = \Omega^\theta\partial_\theta + \Omega^r\partial_r \tag{6}$$

$$\Omega^2_{LT} = g_{rr}(\Omega^r)^2 + g_{\theta\theta}(\Omega^\theta)^2 \tag{7}$$

Next, in the orthonormal 'Copernican' basis at rest in the rotating spacetime, with our choice of polar cooridnates, Ω_{LT} can be written as

$$\vec{\Omega}_{LT} = \sqrt{g_{rr}}\Omega^r\hat{r} + \sqrt{g_{\theta\theta}}\Omega^\theta\hat{\theta} \tag{8}$$

where, $\hat{\theta}$ is the unit vector along the direction θ. Our final result of LT precession in PD spacetime is then,

$$\vec{\Omega}^{PD}_{LT} = \frac{\sqrt{\Delta}}{p}\left[\frac{a(\Xi\cos\theta + (2n - a\cos\theta)\frac{a}{\ell^2}\sin^2\theta)}{\Delta - a^2\Xi\sin^2\theta} - \frac{a\cos\theta - n}{p^2}\right]\hat{r} + \frac{\sqrt{\Xi}}{p}a\sin\theta\left[\frac{r - M - \frac{r}{\ell^2}(a^2 + 2r^2 + 6n^2)}{\Delta - a^2\Xi\sin^2\theta} - \frac{r}{p^2}\right]\hat{\theta} \tag{9}$$

Now, from the above expression we can easily derive the LT precession rates for various axisymmetric stationary spacetimes as the special cases of PD spacetime.

B. Schwarzschild and Schwarzschild-de Sitter spacetimes

As the Schwarzschild and Schwarzschild-de Sitter spacetimes both are static and $a = \Lambda = Q_c = Q_m = n = 0$, the inertial frames are not dragged along it. So, we can't see any LT effect in these spacetimes.

C. Kerr spacetimes

LT precession rate for non-extremal Kerr spacetime is already discussed in details in our paper. Setting $\Lambda = Q_c = Q_m = n = 0$ in eqn.(9) we can recover our result (eqn.(12) of [1]).

D. Kerr-Newman spacetime

Rotating black hole spacetimes with electric charge Q_c and magnetic charge Q is described by Kerr-Newman metric, which is very important in General Relativity. Setting $\Lambda = n = 0$ in eqn.(9), we can easily get the LT precession in Kerr-Newman spacetime. It is thus following (taking $Q_c^2 + Q_m^2 = Q^2$),

$$\vec{\Omega}^{KN}_{LT} = \frac{a}{\rho^3(\rho^2 - 2Mr + Q^2)}\left[\sqrt{\Delta}(2Mr - Q^2)\cos\theta\hat{r} + (M(2r^2 - \rho^2) + rQ^2)\sin\theta\hat{\theta}\right] \tag{10}$$

In the Kerr-Newman spacetime,

$$\Delta = r^2 - 2Mr + a^2 + Q^2 \tag{11}$$

and,

$$\rho^2 = r^2 + a^2\cos^2\theta \tag{12}$$

From the expression (10) of LT precession in Kerr-Newman spacetime we can see that at the polar plane, the LT precession vanishes for the orbit $r = \frac{Q^2}{2M}$, though the spacetime is rotating ($a \neq 0$).

E. Kerr-de Sitter spacetimes

Kerr-de Sitter spacetime is more realistic, when we do not neglect the Cosmological constant parameter (though its value is very small, it may be very useful in some cases, where we need to very precise calculation). Setting $n = Q_c = Q_m = 0$, we get the following expression for Kerr-de Sitter spacetime

$$\vec{\Omega}_{LT}^{KdS} = \frac{a}{\rho^3(\rho^2 - 2Mr - \frac{1}{\ell^2}(a^4 + a^2r^2 - a^4\sin^2\theta\cos^2\theta))}\Big[\sqrt{\Delta}(2Mr + \frac{1}{\ell^2}\rho^4)\cos\theta\hat{r}$$
$$+\sqrt{\Xi}[M(2r^2 - \rho^2) + \frac{r}{\ell^2}\{a^4 + a^2r^2 - a^4\sin^2\theta\cos^2\theta - \rho^2(a^2 + 2r^2)\}])\sin\theta\hat{\theta}\Big] \tag{13}$$

Where,

$$\Delta = (r^2 + a^2)(1 - \frac{r^2}{\ell^2}) - 2Mr \tag{14}$$

and

$$\Xi = 1 + \frac{a^2}{\ell^2}\cos^2\theta \tag{15}$$

F. Kerr-Taub-NUT spacetime

If we set $Q_c = Q_m = \Lambda = 0$ in eqn. (9), we get

$$\vec{\Omega}_{LT}^{KTN} = \frac{\sqrt{\Delta}}{p}\left[\frac{a\cos\theta}{\rho^2 - 2Mr - n^2} - \frac{a\cos\theta - n}{p^2}\right]\hat{r}$$
$$+ \frac{a\sin\theta}{p}\left[\frac{r - M}{\rho^2 - 2Mr - n^2} - \frac{r}{p^2}\right]\hat{\theta} \tag{16}$$

Where, $\Delta = r^2 + a^2 - n^2 - 2Mr$
and, $p^2 = r^2 + (n - a\cos\theta)^2$. The above solutions other than Kerr-Taub-NUT spacetime are not very interesting. These are as usual like Kerr spacetime. The source of the LT precession is Kerr parameter a. If $a = 0$, the LT precession vanishes. But, more surprising thing is that when $a = 0$ in Kerr-Taub-NUT spacetime, the LT precession does not vanish. This means that though the angular momentum (J) of this spacetime is vanished, the inertial frames are dragged along it with the following precession rate

$$\vec{\Omega}_{LT}^{TN} = \frac{n\sqrt{\Delta}}{p^3}\hat{r} \tag{17}$$

We can express it in more general, we mean we can check the precession rate in the case of Plebański-Demiański (PD) spacetimes with vanishing Kerr parameter ($a = 0$). In this case, the expression of LT precession rate is same as eqn. (17), but, in this case,

$$\Delta = (r^2 - n^2)(1 - \frac{1}{\ell^2}(r^2 + 3n^2))$$
$$- 2Mr + Q_c^2 + Q_m^2 - 4\frac{1}{\ell^2}n^2r^2$$
$$p^2 = r^2 + n^2 \tag{18}$$

We can also see that the precession rate is independent of θ. From the above eqn.(17), we can easily see that the LT precession vanishes if the NUT charge vanishes. Because, if $n = 0$, the Taub-NUT spacetime reduces to static Schwarzschild spacetime. So, we should study about this precession in some details.

G. NUT spacetime

The NUT spacetime is a geometrically stationary, spherically symmetric vacuum solution of Einstein equation with NUT charge (n). The Einstein-Hilbert action requires no modification to accommodate this

NUT charge or "dual mass". This dual mass is an intrinsic feature of general relativity. If the NUT charge vanishes, the metric reduces to Schwarzschild spacetime. There are four linearly independent Killing vectors $\xi_t, \xi_x, \xi_y, \xi_z$ which satisfy the same commutation relations as in the Scwarzschild case,

$$[\xi_a, \xi_b] = -\varepsilon_{abc}\xi_c, \tag{19}$$

where a, b, c has the values x, y, z and, also for stationarity condition we must have

$$[\xi_t, \xi_a] = 0 \tag{20}$$

However, let us first consider the line element (of NUT spacetime), which is presented by Newman et. al.

$$ds^2 = -f(r)\left[dt + 4n\sin^2\frac{\theta}{2}d\phi\right]^2 + \frac{1}{f(r)}dr^2 + (r^2+n^2)(d\theta^2 + \sin^2\theta d\phi^2) \tag{21}$$

where,

$$f(r) = \frac{r^2 - 2Mr - n^2}{r^2+n^2} \tag{22}$$

Here, M represents the "gravitoelectric mass" or 'mass' and n represents the "gravitomagnetic mass" or 'dual mass' of this spacetimes. The metric satisfies the Einstein field equations in vacuum

$$G_{\mu\nu} = R_{\mu\nu} = 0 \tag{23}$$

For $f(r) = 0$, there are co-ordinate singularities at

$$r = r_{\pm} = M \pm \sqrt{M^2+n^2} \tag{24}$$

at NUT spacetime and $r_{\pm}$ correspond to *horizons* in this geometry. Actually, they are horizons in this sence that $f(r)$ changes the sign from positive to negative across the horizon and r changes from spacelike to timelike. So, only for stationary axisymmetric NUT spacetime, the precession rate (which could be easily got from eqn.(17) for $Q_c = Q_m = \Lambda = 0$) will be

$$\vec{\Omega}_{LT}^{TN} = \frac{n(r^2 - 2Mr - n^2)^{\frac{1}{2}}}{(r^2+n^2)^{\frac{3}{2}}}\hat{r} \tag{25}$$

If n is vanished, Ω is also vanished. Again, there is also an another case when the LT precession rate vanishes (though $n \neq 0$), that is happen at the points of co-ordinate singularities ($r = r_{\pm}$). Otherwise, LT precession will be continued in that spacetime though the intrinsic *angular momentum*(J) of the spacetime is *zero*. We already know that LT precession could not be observed in Schwarzschild geometry which is a spherically symmetric spacetimes. But, Taub-NUT spacetime is a spherically symmetric spacetime and we can observe LT precession, here. So, what the origin of this precession is. As the massless spacetime is also well-defined (see, for example, appendix of [9]), we can also write down the precession rate only for massless dual mass (NUT charge n can be regarded as dual mass) solutions of NUT spacetime. It would be

$$\vec{\Omega}_{LT}^{TN} = \frac{n(r^2 - n^2)^{\frac{1}{2}}}{(r^2+n^2)^{\frac{3}{2}}}\hat{r} \tag{26}$$



FIG. 1. Plot of r vs Ω_{LT}^{TN} for $l = 3$ m

At, the points of singularities ($r = \pm n$), the LT precession is vanished same as the previous case. We can visualize this characteristics with a graph(FIG. 1). In this figure, we have plotted precession rate (Ω_{LT}^{TN}) along the y axis and distance (r) along the x axis. Here, we observe that for $n = 3$, the LT precession is starting for $r > 3$ and continued to the infinity. Setting $\frac{d\Omega_{LT}^{TN}}{dr} = 0$, we get that Ω_{LT}^{TN} is maximum at $r = \sqrt{2}n$. In our figure, this value is $r = 3\sqrt{2} = 4.24$ m. Now, we are not interested for $r < 3$. Our formulas are not comfortable in that regions. From our precession rate formulas (17,25,26) at "dual mass" spacetimes we can see that the precession rate (Ω_{LT}^{TN}) is the same, starting from the polar region to the equatorial plane for a fixed distance. Ω_{LT}^{TN} depends only on distance (r) of the test gyroscope from the 'dual mass'. So, we conclude that the source of LT precession in NUT spacetime is just the "dual mass". So, the *"dual mass"* is same as the *"angular momentum monopole"*, in NUT spacetime.

III. EXTREMAL CASE

A. Extremal Plebański-Demiański Spacetime

In this section, we would like to describe the LT precession in extremal Plebański-Demiański spacetime, whose non-extremal case is already described in the previous section. To get the extremality condition in PD spacetimes we should first determine the horizon r_h is given by $\Delta|_{r=r_h} = 0$. We can make a comparison of coefficients in

$$\begin{aligned}\Delta &= -\frac{1}{\ell^2}r^4 + (1 - \frac{a^2}{\ell^2} - \frac{6n^2}{\ell^2})r^2 - 2Mr \\ &\quad + \left[(a^2 - n^2)(1 - \frac{3n^2}{\ell^2}) + Q_c^2 + Q_m^2\right] \\ &= -\frac{1}{\ell^2}[r^4 + (a^2 + 6n^2 - \ell^2)r^2 + 2M\ell^2 r + b] \\ &= -\frac{1}{\ell^2}\Pi_{i=1}^4 (r - r_{hi}) \end{aligned} \tag{27}$$

where,

$$b = (a^2 - n^2)(3n^2 - \ell^2) - \ell^2(Q_c^2 + Q_m^2) \tag{28}$$

and r_{hi}(i = 1, 2, 3, 4) denotes the zeros of Δ[5]. From this comparison we can conclude that for the PD (when Λ is negative, "anti-de-Sitter" or "AdS") black hole, there are two separated positive horizons at most, and Δ is positive outside the outer horizon of the PD black hole. In the same way, we can conclude for the PD (when Λ is positive, "de-Sitter" or "dS") black hole that there are three separated positive horizons at most, and Δ is negative outside the outer horizon of the black hole. Both of the above cases, the when two horizons of the PD black hole coincide, the black hole is extremal [15].

If we consider the extremal PD black hole, we have to make a comparison of coefficients in

$$\begin{aligned}\Delta &= (r - x)^2(a_2 r^2 + a_1 r + a_0) \\ &= -\frac{1}{\ell^2}[r^4 + (a^2 + 6n^2 - \ell^2)r^2 + 2Mr\ell^2 + b]\end{aligned} \tag{29}$$

with a_0, a_1, a_2 being real[5]. From this comparison we can get the following for PD ("AdS") spacetime,

$$\frac{b_A}{x^2} - 3x^2 = a^2 + 6n^2 + \ell^2 \tag{30}$$

$$x^3 - \frac{b_A}{x} = -M\ell^2 \tag{31}$$

where, b_A represents the value of b at PD ("AdS") spacetimes.

$$b_A = (a^2 - n^2)(3n^2 + \ell^2) + \ell^2(Q_c^2 + Q_m^2) \tag{32}$$

Solving equation (30) for x, we get

$$x = \sqrt{\frac{1}{6}\left[-(\ell^2 + a^2 + 6n^2) + \sqrt{(\ell^2 + a^2 + 6n^2)^2 + 12b_A}\right]} \tag{33}$$

Similarly, we can obtain for PD("dS") black hole ,

$$\frac{b}{x^2} - 3x^2 = a^2 + 6n^2 - \ell^2 \tag{34}$$

$$x^3 - \frac{b}{x} = M\ell^2 \tag{35}$$

In these equation x is positive and related to the coincided horizon of the extremal PD (for "dS") black hole.

Solving equation (34) for x, we get

$$x_+ = \sqrt{\frac{1}{6}\left[\ell^2 - a^2 - 6n^2 + \sqrt{(\ell^2 - a^2 - 6n^2)^2 + 12b}\right]} \tag{36}$$

$$x_- = \sqrt{\frac{1}{6}\left[\ell^2 - a^2 - 6n^2 - \sqrt{(\ell^2 - a^2 - 6n^2)^2 + 12b}\right]} \tag{37}$$

where, x_+ and x_- indicate the outer horizon and inner horizon, respectively. This can be seen by calculating

$$\frac{d^2\Delta}{dr^2}|_{r=x_+} = -\frac{2}{\ell^2}\sqrt{(\ell^2 - a^2 - 6n^2)^2 + 12b} \tag{38}$$

and

$$\frac{d^2\Delta}{dr^2}|_{r=x_-} = \frac{2}{\ell^2}\sqrt{(\ell^2 - a^2 - 6n^2)^2 + 12b} \tag{39}$$

For the PD("dS") black hole, on the outer extremal horizon, $\frac{d\Delta}{dr} = 0$ and $\frac{d^2\Delta}{dr^2} < 0$ and on the inner extremal horizon $\frac{d^2\Delta}{dr^2} > 0$. Now, we can solve M and a from the two equations(34, 35).

$$M = \frac{x\left[x^4 + 2x^2(3n^2 - \ell^2) + (3n^2 - \ell^2)(7n^2 - \ell^2) + \ell^2(Q_c^2 + Q_m^2)\right]}{\ell^2(\ell^2 + x^2 - 3n^2)} \tag{40}$$

$$a_e^2 = \frac{3x^4 + (6n^2 - \ell^2)x^2 + n^2(3n^2 - \ell^2) + \ell^2(Q_c^2 + Q_m^2)}{(3n^2 - \ell^2 - x^2)} \tag{41}$$

From the above values of a_e^2 and M, we get

$$a_e^2 = -M\ell^2 \frac{\left[3x^4 + (6n^2 - \ell^2)x^2 + n^2(3n^2 - \ell^2) + \ell^2(Q_c^2 + Q_m^2)\right]}{x\left[x^4 + 2x^2(3n^2 - \ell^2) + (3n^2 - \ell^2)(7n^2 - \ell^2) + \ell^2(Q_c^2 + Q_m^2)\right]} \tag{42}$$

The ranges of x and a_e are determined from the following expressions

$$x^2 < \left(\frac{\ell^2 + \ell\sqrt{\ell^2 - 12Q^2}}{6} - n^2\right) \tag{43}$$

$$0 < a_e^2 < \left[(7\ell^2 - 24n^2) - \sqrt{(7\ell^2 - 24n^2)^2 - (\ell^4 - 12\ell^2(Q_c^2 + Q_m^2))}\right] \tag{44}$$

Due to the presence of Cosmological constant, there exist four roots of x in eqn. (41). When Cosmological constant $\frac{1}{l^2} \to 0$, eqn.(40) and eqn.(41) reduces to

$$x = M \tag{45}$$

and,

$$a_e^2 = x^2 + n^2 - Q_c^2 - Q_m^2 \tag{46}$$

or,

$$a_e^2 = M^2 + n^2 - Q_c^2 - Q_m^2 \tag{47}$$

respectively.

Now, the line element of extremal PD spacetimes can be written as,

$$ds^2 = -\frac{\Delta_e}{p_e^2}(dt - A_e d\phi)^2 + \frac{p_e^2}{\Delta_e}dr^2 + \frac{p_e^2}{\Xi_e}d\theta^2 + \frac{\Xi_e}{p_e^2}\sin^2\theta(a_e dt - B_e d\phi)^2 \tag{48}$$

and, the final LT precession rate in extremal PD spacetime is,

$$\begin{aligned}\vec{\Omega}_{LT}^{ePD} = \frac{\sqrt{\Delta_e}}{p_e}&\left[\frac{a_e(\Xi_e\cos\theta + (2n - a_e\cos\theta)\frac{a_e}{\ell^2}\sin^2\theta)}{\Delta_e - a_e^2\Xi_e\sin^2\theta} - \frac{a_e\cos\theta - n}{p_e^2}\right]\hat{r}\\ &+ \frac{\sqrt{\Xi_e}}{p_e}a_e\sin\theta\left[\frac{r - M - \frac{r}{\ell^2}(a_e^2 + 2r^2 + 6n^2)}{\Delta_e - a_e^2\Xi_e\sin^2\theta} - \frac{r}{p_e^2}\right]\hat{\theta}\end{aligned} \tag{49}$$

where,

$$\begin{aligned}\Delta_e &= -\frac{1}{\ell^2}(r - x)^2\left(r^2 + 2rx + \frac{b}{x^2}\right)\\ \Xi_e &= 1 + \frac{a_e^2}{\ell^2}\cos^2\theta - \frac{4a_e n}{\ell^2}\cos\theta\\ p_e &= r^2 + (n - a_e\cos\theta)^2\\ A_e &= a_e\sin^2\theta + 2n\cos\theta\\ B_e &= r^2 + a_e^2 + n^2\end{aligned} \tag{50}$$

and, the value of a_e is determined from the eqn.(41) and the range of x and a_e ('e' stands for the *extremal* case) are determined from the eqn. (43) and (44), respectively. It could be noted that, for the extremal PD("dS"), spacetimes, *there are upper limiting values for angular momentum and extremal horizon of the black hole.* Substituting all the above mentioned values and ranges in eqn.(49), we get the exact LT precession rate in extremal PD spacetimes.

B. Extremal Kerr Spacetime

Substituting $a_e = M$ in equation (12) of [1]) or substituting $a = M$ and $\Lambda = Q_c = Q_m = n = 0$ in equation (49) we can easily get the LT precession rate in extremal Kerr spacetime.

$$\Omega_{LT}^{eK}(r,\theta) = \frac{M^2}{(r^2 + M^2\cos^2\theta)^{\frac{3}{2}}(r^2 - 2Mr + M^2\cos^2\theta)}\left[4(r - M)^2r^2\cos^2\theta + (r^2 - M^2\cos^2\theta)^2\sin^2\theta\right]^{\frac{1}{2}} \tag{51}$$

The above result is also coming from the eqn.(49), the general LT precession rate for extremal PD spacetime.
Case I: On the polar region i.e. $\theta = 0$ the Ω_{LT} becomes

$$\Omega^{eK}_{LT} = \frac{2M^2 r}{(r^2+M^2)^{3/2}(r-M)} \tag{52}$$

Case II: On the equator i.e. $\theta = \pi/2$, the Ω_{LT} becomes

$$\Omega^{eK}_{LT} = \frac{M^2}{r^2(r-2M)} \tag{53}$$

It could be easily seen that Ω_{LT} is diverges at $r = M$. Since $r = M$ is the only direct ISCO in extremal Kerr geometry which coincides with the principal null geodesic generator of the horizon[16, 17] and it is also the radius of the horizon which is a null surface. Our general LT precession formula is derived only considering that the observer is rest in a timelike Killing vector field. We have not incorporate the LT effect for any null geodesics. So, our formula is valid for $r > M$ (outside the horizon).

C. Extremal Kerr-Newman Spacetime

Substituting $Q_c^2 + Q_m^2 = M^2 - a_e^2$ and $\Lambda = n = 0$ in equation (9), we can obtain the following LT precession rate for extremal KN blackhole

$$\begin{aligned}\vec{\Omega}^{eKN}_{LT} = & \frac{a_e}{\rho^3[(r-M)^2 - a_e^2\sin^2\theta]}\Big[(r-M)(a_e^2 - M^2 + 2Mr)\cos\theta\hat{r} \\ & +\{r(a_e^2 - M^2 + 2Mr)\} - M\rho^2)\sin\theta\hat{\theta}\Big]\end{aligned} \tag{54}$$

where

$$\rho^2 = r^2 + a_e^2\cos^2\theta \tag{55}$$

The same expression (eqn.54) could be derived easily from eqn.(49).

D. Extremal Kerr-de Sitter Spacetime

Extremal Kerr-de Sitter spacetime is interesting because it involves the Cosmological constant. Setting $n = Q_c = Q_m = 0$ in eqn.(9), we can find the following expression of LT precession at extremal Kerr-de Sitter spacetimes

$$\begin{aligned}\vec{\Omega}^{eKdS}_{LT} = & \frac{a_e}{\rho^3(\rho^2 - 2Mr - \frac{1}{\ell^2}(a_e^4 + a_e^2 r^2 - a_e^4\sin^2\theta\cos^2\theta))}\Big[\sqrt{\Delta_e}(2Mr + \frac{\rho^4}{\ell^2})\cos\theta\hat{r} \\ & +\sqrt{\Xi_e}[M(2r^2-\rho^2) + \frac{r}{\ell^2}\{a_e^4 + a_e^2 r^2 - a_e^4\sin^2\theta\cos^2\theta - \rho^2(a_e^2+2r^2)\}])\sin\theta\hat{\theta}\Big]\end{aligned} \tag{56}$$

where,

$$\begin{aligned}\rho^2 &= r^2 + a_e^2\cos^2\theta \\ \Delta_e &= -\frac{1}{\ell^2}(r-x)^2\left(r^2 + 2xr - \frac{a^2\ell^2}{x^2}\right) \\ \Xi_e &= 1 + \frac{a_e^2}{\ell^2}\cos^2\theta\end{aligned} \tag{57}$$

Using equations (41, 40), we obtain the value for a and M are

$$\begin{aligned}a_e^2 &= \frac{(\ell^2 - 3x^2)x^2}{(\ell^2 + x^2)} \\ M &= \frac{x(\ell^2 - x^2)^2}{\ell^2(\ell^2 + x^2)}\end{aligned} \tag{58}$$

where the horizons are at

$$\begin{aligned}x_+ &= \sqrt{\frac{1}{6}\left[\ell^2 - a^2 + \sqrt{(\ell^2-a^2)^2 - 12a^2\ell^2}\right]} \\ x_- &= \sqrt{\frac{1}{6}\left[\ell^2 - a^2 - \sqrt{(\ell^2-a^2)^2 - 12a^2\ell^2}\right]}\end{aligned} \tag{59}$$

where, x_+ and x_- indicate the outer horizon and inner horizon. The range of a_e^2 is $0 < a_e^2 < (7-4\sqrt{3})\ell^2$ and $x^2 < \ell^2/3$, which is already discussed in [15]. Substituting all the above values in eqn.(56) and taking the ranges of a_e and x, we get the exact LT precession rate in extremal Kerr-dS spacetimes.

E. Extremal Kerr-Taub-NUT spacetime

To derive the extremal limit in Kerr-Taub-NUT spacetime we set,

$$\Delta = r^2 - 2Mr + a^2 - n^2 = 0 \tag{60}$$

Solving for r, we get two horizons which are located at $r_\pm = M \pm \sqrt{M^2 + n^2 - a^2}$. So, we get the extremality condition $(r_+ = r_-)$ for Kerr-Taub-NUT spacetimes is $a_e^2 = M^2 + n^2$. If we set $Q_c = Q_m = \Lambda = 0$ and $M^2 + n^2 = a_e^2$ in eqn. (9), we get the following exact LT precession rate at extremal Kerr-Taub-NUT spacetime.

$$\vec{\Omega}_{LT}^{eKTN} = \frac{(r-M)}{p}\left[\frac{a_e\cos\theta}{(r-M)^2 - a_e^2\sin^2\theta} - \frac{a_e\cos\theta \mp \sqrt{a_e^2 - M^2}}{p^2}\right]\hat{r} + \frac{a_e\sin\theta}{p}\left[\frac{r-M}{(r-M)^2 - a_e^2\sin^2\theta} - \frac{r}{p^2}\right]\hat{\theta} \tag{61}$$

Where $p^2 = r^2 + (\pm\sqrt{a_e^2 - M^2} - a_e\cos\theta)^2$.

IV. DISCUSSIONS

In this work we have explicitly derived the Lense Thirring precession frequency for extremal and non-extremal Plebański-Demiański(PD) spacetime. The PD family of solutions are the solutions of Einstein field equations which contain a number of well known blackhole solutions. Among them including Schwarzschild, Schwarzschild-de Sitter, Kerr, Kerr-ads, Kerr-Newman, Kerr-Taub-NUT etc. We observe that LT precession frequency strongly depends upon the different parameters like mass M, spin a, Cosmological constant Λ, NUT charge n, electric charge Q_c and magnetic charge Q_m. An interesting point we have seen that for NUT parameter LT precession occurs solely due to the “dual mass”. This “dual mass” is equivalent to angular momentum monopole (n) in NUT spacetime. For our completeness we have also deduced the LT precession for extremal PD spacetime and also for others axisymmetric PD-like spacetimes. To get the LT precession rate in extremal Kerr-Taub-NUT-de-Sitter spacetime, the basic procedure is same as the PD spacetimes with the additional requirement is $Q_c^2 + Q_m^2 = 0$ in equations (40) and (41) and the range of $x^2 < (\frac{\ell^2}{3} - n^2)$ and a_e^2 has a range of $0 < a_e^2 < (7\ell^2 - 24n^2) - 4\sqrt{3(\ell^2 - 3n^2)(\ell^2 - 4n^2)}$. As there is no any valid extremality condition at NUT spacetime (with M or without M), we could not get any *real* LT precession rate due to frame-dragging effect. Since, the direct ISCO coincides with the principal null geodesic generator[16, 17] in extremal Kerr spacetime, we are unable to discuss the LT precession at that particular geodesic. So this formula is not valid for the domain of $r \leq M$ for the extremal Kerr spacetime. Here, our formula is valid only for $r > M$. In general, our most general formula for LT precession in stationary spacetime is valid only outside the horizon, as our observer is in timelike Killing vector field. Our formula is not valid on the horizon and inside the horizon. We will discuss about this problem in near future.

Acknowledgments : We would like to thank Prof. Parthasarathi Majumdar for his encouragement and guidance, besides the long blackboard discussions on this topic.

[1] Chakraborty, C., Majumdar, P., arXiv:1205.6722 [gr-qc].
[2] Lense, J., Thirring, H., *Phys. Z.* **19**, 156-163, (1918).
[3] Kerr, R.P., *Phys. Rev. Lett.*,**11**, 237-238, (1963).
[4] Plebański, J., Demiański, M.,*Ann. Phys.* (N.Y.)**98**, 98(1976)
[5] Hackmann, E., Lammerzahl, C., *Phys. Rev.D*,**85**, 044049 (2012).
[6] Misner, C. W., *J. Math. Phys.*, **4**, 924 (1963).
[7] Taub, A.H.,*Ann. of Maths.*, **53**, 3 (1951).
[8] Newman, E., Tamburino, L., Unti, T.,*J. Math. Phys.*, **4**, 7 (1963).
[9] Ramaswamy, S., Sen, A., *J. Math. Phys. (N.Y.)* **22**, 2612 (1981).
[10] Ramaswamy, S., Sen, A., *Phys. Rev. Lett.* **57**, 8 (1986).
[11] Lynden-Bell1, D., Nouri-Zonoz, M.,arXiv:9612049v1 [gr-qc].
[12] Mueller, M., Perry, M.J.,*Class. Quantum Grav.* **3**, 65-69 (1986).
[13] Chandrasekhar, S., *The Mathematical Theory of Black Holes*, Oxford (1992).
[14] Hartle, J. B., *Gravity: An introduction to Einstein's General relativity*, Pearson (2009).
[15] Li, Y. et. al., *Class. Quantum Grav.* **28**, 225006 (2011).
[16] Pradhan, P., Majumdar, P., arXiv:1108.2333v1 [gr-qc].
[17] Pradhan, P., (in preparation).